



ISSN 2307-8235 (online)


IUCN 2008: T132725A100453919

Scope: Global

Language: English

Cephalopholis aurantia, Golden Hind

Assessment by: Amorim, P., Das, D., Rhodes, K., Russell, B., Law, C., Suharti, S., Samoilys, M., Nair, R., Ma, K., To, A., Myers, R. & Choat, J.H.



View on www.iucnredlist.org

Citation: Amorim, P., Das, D., Rhodes, K., Russell, B., Law, C., Suharti, S., Samoilys, M., Nair, R., Ma, K., To, A., Myers, R. & Choat, J.H. 2018. *Cephalopholis aurantia*. The IUCN Red List of Threatened Species 2018: e.T132725A100453919. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T132725A100453919.en>

Copyright: © 2018 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Actinopterygii	Perciformes	Epinephelidae

Taxon Name: *Cephalopholis aurantia* (Valenciennes, 1828)

Synonym(s):

- *Bodianus indelibilis* Fowler, 1904
- *Cephalopholis analis* (Valenciennes, 1828)
- *Cephalopholis aurantius* (Valenciennes, 1828)
- *Cephalopholis indelibilis* (Fowler, 1904)
- *Cephalopholis obtusauris* Evermann & Seale, 1907
- *Cephalopholis obtusaurus* Evermann & Seale, 1907
- *Epinephelus analis* (Valenciennes, 1828)
- *Epinephelus aurantius* (Valenciennes, 1828)
- *Epinephelus miltostigma* Bleeker, 1873
- *Serranus analis* Valenciennes, 1828
- *Serranus aurantius* Valenciennes, 1828
- *Serranus rufus* Hombron & Jacquinot, 1853

Common Name(s):

- English: Golden Hind, Golden Rockcod, Orange Cod
- French: Cuisinier, Maconde, Mamam Rouge, Rouge Batard, Rouge Peau Dure, Vieille Doree
- Spanish: Cherna Dorada

Assessment Information

Red List Category & Criteria: Least Concern [ver 3.1](#)

Year Published: 2018

Date Assessed: November 18, 2016

Justification:

Cephalopholis aurantia is a widely distributed and deep-living species that is infrequently captured. There are no known major threats; therefore, it is listed as Least Concern. The change in status from the previous assessment reflects an improved application of the Red List categories and criteria, as well as a better understanding of available data.

Previously Published Red List Assessments

2008 – Data Deficient (DD)

<http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T132725A3433665.en>

Geographic Range

Range Description:

This Indo-Pacific species is distributed in the islands of the western Indian Ocean to Japan and the

central Pacific. It is also recorded from the Andaman and Nicobar Islands (Sachithanandan and Mohan 2014), from Rowley Shoals in Australian waters and Christmas Island in the Indian Ocean (Bray 2017) and Tanega-shima Island in Kagoshima, Japan (Koeda and Motomura 2015). The species is also recorded around Howland Island in the Pacific (B. Mundy pers. comm.). A single specimen has been caught off Natal, South Africa. Heemstra and Randall (1993) reported no other confirmed records from other continental localities of East Africa. There is one unconfirmed record from Mozambique (S. Fennessy pers. comm. 2016) and it has been reported by fishermen in Mafia Island, Tanzania (Gaspere *et al.* 2015). Its occurrence in Malaysia is questionable. It is recorded from Cochin, India, but the actual provenance of the specimens are unknown (R. Nair pers. comm. 2016).

The specific localities in which it occurs are as follows: American Samoa, Australia, China, South China Sea (Spratly Islands), Taiwan, Christmas Island, Cook Islands, Djibouti, Fiji, French Polynesia (Tahiti, Society Islands, Tuamotu Islands), Guam, India (Lakshadweep and Andaman and Nicobar), Indonesia (Bali, Lesser Sunda Islands, Moluccas, Papua, Sulawesi, Sumatra), Japan (Ryukyu Islands), Madagascar, Maldives, Marshall Islands, Mauritius (Rodrigues), Micronesia, Mozambique, New Caledonia (not in Loyalty Islands), Northern Mariana Islands, Palau, Papua New Guinea, Philippines, Pitcairn, Réunion, Samoa, Seychelles, Solomon Islands, South Africa (KwaZulu-Natal), Thailand (not shown in map), Tonga, Tuvalu.

Country Occurrence:



Native: American Samoa; Australia; China; Christmas Island; Comoros; Disputed Territory (Spratly Is.); Fiji; French Polynesia; French Southern Territories (Mozambique Channel Is.); Guam; India (Andaman Is., Laccadive Is., Nicobar Is.); Indonesia; Japan; Kenya; Kiribati (Phoenix Is.); Madagascar; Malaysia; Maldives; Marshall Islands; Mauritius (Mauritius (main island), Rodrigues); Mayotte; Micronesia, Federated States of ; Mozambique; Myanmar; New Caledonia; Niue; Northern Mariana Islands; Palau; Papua New Guinea; Philippines; Réunion; Samoa; Seychelles; Solomon Islands; Somalia; South Africa; Taiwan, Province of China; Tanzania, United Republic of; Thailand; Timor-Leste; Tonga; Tuvalu; Vanuatu; Wallis and Futuna; Yemen

FAO Marine Fishing Areas:

Native: Indian Ocean - western, Indian Ocean - eastern, Pacific - western central, Pacific - northwest

Distribution Map

Cephalopholis aurantia




Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community

Range

Extant (resident)

Compiled by:

IUCN Grouper and Wrasse Specialist Group



Population

This grouper is relatively poorly known due to the deep depths which it inhabits; therefore, population data are limited. At times, it can be misidentified as *Cephalopholis sonnerati*.

Current Population Trend: Unknown

Habitat and Ecology (see Appendix for additional information)

This species inhabits steep seaward reefs, usually at depths of 100 to 250 m. It has been found at depths as shallow as 20 m in Australia (Bray 2017), but this may be outside its natural depth range. Its maximum total length is 30 cm (Craig *et al.* 2011), and a report of 60 cm is considered invalid (Heemstra and Randall 1993). A hybrid specimen of this species between *Cephalopholis aurantia* and *C. spiloparaea* has been found on a barrier reef in New Caledonia at 50-60 m depth (Randall and Justine 2008).

Systems: Marine

Use and Trade

This species is occasionally incidentally caught in deep line fisheries, and is not targeted by commercial fisheries. It has been recorded in landings in the Andaman archipelago, India (Sachithanandam and Mohan 2014). In Mafia Island, Tanzania, fishermen interviewed on grouper fishing reported this species is caught infrequently compared to other species (Gaspere *et al.* 2015). It has not been recorded in Indonesia deep water fishery monitoring surveys (P. Mous and J. Pet, The Nature Conservancy Indonesia Fisheries Conservation Program database, unpublished, accessed 2016). It has also not been recorded from Solomon Island market surveys during which 65,000 specimens were investigated (Solomon Islands Ministry of Fisheries and Marine Resources unpublished database, accessed 2016).

Threats (see Appendix for additional information)

There are no known major threats.

Conservation Actions (see Appendix for additional information)

There are no species-specific conservation measures.

Credits

Assessor(s): Amorim, P., Das, D., Rhodes, K., Russell, B., Law, C., Suharti, S., Samoilys, M., Nair, R., Ma, K., To, A., Myers, R. & Choat, J.H.

Reviewer(s): Linardich, C.

Facilitators(s) and Carpenter, K.E.

Compiler(s):

Bibliography

Bray, D.J. 2017. *Cephalopholis aurantia*. *Fishes of Australia*
<http://fishesofaustralia.net.au/home/species/5173>.

Craig, M.T., Sadovy de Mitcheson, Y.J. and Heemstra, P.C. 2011. *Groupers of the World: A Field and Market Guide*. NISC Press, South Africa.

Gaspere, L., Bryceson, I. and Kulindwa, K. 2015. Complementarity of fishers' traditional ecological knowledge and conventional science: Contributions to the management of groupers (Epinephelinae) fisheries around Mafia Island, Tanzania. *Ocean and Coastal Management* 114: 88-101.

Heemstra, P.C. and Randall, J.E. 1993. *FAO species catalogue. Vol. 16. Groupers of the world (Family Serranidae, Subfamily Epinephelinae). An annotated and illustrated catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date*. FAO, Rome.

Hoese, D.F., Bray, D.J., Paxton, J.R. and Allen, G.R. 2006. Fishes. In: P.L. Beasley and A. Wells (eds), *Zoological Catalogue of Australia*, pp. 2178. Australian Biological Resources Study and Australia's Commonwealth Scientific and Industrial Research Organisation (CSIRO), Australia.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-2. Available at: www.iucnredlist.org. (Accessed: 15 November 2018).

Koeda, K., Motomura, H. 2015. First record of *Cephalopholis aurantia* (Perciformes: Serranidae) from Tanega-shima island and the Tokara Islands, Kagoshima, Japan. *Nature of Kagoshima* 41(47-52).

Randall, J.E. and Justine, J. 2008. *Cephalopholis aurantia* × *C. spiloparaea*, a Hybrid Serranid Fish from New Caledonia. *The Raffles Bulletin of Zoology* 56(1): 157-159.

Sachithanandam, V. and Mohan, P.M. 2014. New Distribution Record of *Cephalopholis aurantia* Valenciennes, 1828 (Pisces:Serranidae) Golden Hind to Indian Waters from Andaman and Nicobar Islands. *Indian Journal of Geo-Marine Sciences* 43(12): 2233-2235.

The Nature Conservancy and Worldwide Fund for Nature. 2003. Report on a rapid ecological assessment of the Raja Ampat Islands, Papua, eastern Indonesia.

Citation

Amorim, P., Das, D., Rhodes, K., Russell, B., Law, C., Suharti, S., Samoilys, M., Nair, R., Ma, K., To, A., Myers, R. & Choat, J.H. 2018. *Cephalopholis aurantia*. The IUCN Red List of Threatened Species 2018: e.T132725A100453919. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T132725A100453919.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the Red List website.

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
9. Marine Neritic -> 9.2. Marine Neritic - Subtidal Rock and Rocky Reefs	Resident	Suitable	Yes

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
5. Biological resource use -> 5.4. Fishing & harvesting aquatic resources -> 5.4.3. Unintentional effects: (subsistence/small scale) [harvest]	Ongoing	Minority (50%)	Unknown	Unknown
	Stresses:	2. Species Stresses -> 2.1. Species mortality		
5. Biological resource use -> 5.4. Fishing & harvesting aquatic resources -> 5.4.4. Unintentional effects: (large scale) [harvest]	Ongoing	Minority (50%)	Unknown	Unknown
	Stresses:	2. Species Stresses -> 2.1. Species mortality		

Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions in Place
In-Place Land/Water Protection and Management
Occur in at least one PA: Unknown

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.3. Life history & ecology

Additional Data Fields

Distribution
Lower depth limit (m): 250
Upper depth limit (m): 100

Population
Population severely fragmented: No

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).